

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark OfficeAttorney Docket No.: STRATA-
06948

Serial No.: 10/087,388

INFORMATION DISCLOSURE STATEMENT BY APPLICANT
(37 CFR § 1.98(b))Applicant: Allen Comer *et al.*

Filing Date: 03/01/02

Group Art Unit:

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

- | | |
|----|--|
| 1 | Berger <i>et al.</i> , Secreted placental alkaline phosphatase: a powerful new quantitative indicator of gene expression in eukaryotic cells, <i>Gene</i> 66:1-10 (1998) |
| 2 | Jaspers <i>et al.</i> , Arsenite Exposure of Cultured Airway Epithelial Cells Activates κ B-dependent Interleukin-8 Gene Expression in the Absence of Nuclear Factor κ B Nuclear Translocation, <i>J. Biol. Chem.</i> 274:31025-31033 (1999) |
| 3 | Muller <i>et al.</i> , Enhanced expression of IL-8 in normal human keratinocytes and human keratinocyte cell line HaCaT <i>in vitro</i> after stimulation with contact sensitizers, tolerogens and irritants, <i>Exp. Dermatol.</i> 3:298-303 (1994) |
| 4 | Newby <i>et al.</i> , Cytokine Release and Cytotoxicity in Human Keratinocytes and Fibroblasts Induced by Phenols and Sodium Dodecyl Sulfate, <i>Soc. of Investigative Dermatology</i> , 115:292-298 (2000) |
| 5 | Fisher <i>et al.</i> , All-Trans Retinoic Acid Induces Cellular R-tinol-Binding Protein in Human Skin <i>In Vivo</i> , <i>Soc. of Investigative Dermatology</i> , 105:80-86 (1995) |
| 6 | Wilmer <i>et al.</i> , Cytokine Induction in Human Epidermal Keratinocytes Exposed to Contact Irritants and Its Relation to Chemical-Induced Inflammation in Mouse Skin, <i>Soc. of Investigative Dermatology</i> , 102:915-922 (1994) |
| 7 | Roguet, <i>Cell Biology and Toxicology</i> , 15:64-75 (1999) |
| 8 | van Ruissen <i>et al.</i> , Differential Effects of Detergents on Keratinocyte Gene Expression, <i>Soc. of Investigative Dermatology</i> , 110:358-363 (1998) |
| 9 | Terunuma <i>et al.</i> , Cytokine mRNA profiles in cultured human skin component cells exposed to various chemicals: a simulation model of epicutaneous stimuli induced by skin barrier perturbation in comparison with that due to exposure to haptens or irritant, <i>J. of Dermatological Science</i> , 26:85-93 (2001) |
| 10 | Medina <i>et al.</i> , Use of Human Skin Equivalent Appligra [®] for <i>In Vitro</i> Assessment of Cumulative Skin Irritation Potential of Topical Products, <i>Toxicology and Applied Pharmacology</i> , 164:38-45 (2000) |
| 11 | Muller-Decker <i>et al.</i> , Keratinocyte-Derived Proinflammatory Key Mediators and Cell Viability as <i>In Vitro</i> Parameters of Irritancy: A Possible Alternative to the Draize Skin Irritation Test, <i>Toxicology and Applied Pharmacology</i> , 127:99-109 (1994) |
| 12 | Abe <i>et al.</i> , Interleukin-8 Gene Repression by Clarithromycin Is Mediated By The activator Protein-1 Binding Site in Human Bronchial Epithelial Cells, <i>Am. J. Respir. Cell Mol. Biol.</i> , 22:51-60 (2000) |

Examiner: /Leon Lankford Jr/

Date Considered: 01/21/2009

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.